

bulletin board ref.: gr-qc/9403046

Quantum Master Equation of Particle in Gas Environment

Lajos Diósi^(a)

KFKI Research Institute for Particle and Nuclear Physics
H-1525 Budapest 114, POB 49, Hungary

Abstract

The evolution of the reduced density operator ρ of Brownian particle is discussed in single collision approach valid typically in low density gas environments. This is the first succesful derivation of quantum friction caused by *local* environmental interactions. We derive a Lindblad master equation for ρ , whose generators are calculated from differential cross section of a single collision between Brownian and gas particles, respectively. The existence of thermal equilibrium for ρ is proved. Master equations proposed earlier are shown to be particular cases of our one.

Brownian motion is one of the most studied classical problem. It has drawn special attention in the quantum case, too. The irreversible dynamics of quantum particle interacting with a stationary environment (e.g. a thermalized reservoir or a stream of molecules, photons, phonons, e.t.c.) is related with many interesting fields in physics. Nonetheless, the quantum counterpart of the classical Brownian motion has not been casted into standard equations since the quantum case is richer than the classical one, especially quantum friction is not straightforward to include correctly. In this Letter we derive quantum frictional equations arising from *local* environmental interactions typical for a gas environment.

In the classical fenomenological theory, the momentum distribution $\rho(\mathbf{p})$ of a Brownian particle of mass M satisfies the Fokker-Planck equation:

$$\frac{d\rho(\mathbf{p})}{dt} = D_{pp}\nabla^2\rho(\mathbf{p}) + \frac{\eta}{M}\nabla(\mathbf{p}\rho(\mathbf{p})) \quad (1)$$

where D_{pp} is the coefficient of momentum diffusion and η is the friction constant. In particular, $D_{pp} = k_B T \eta$ if the environment is in equilibrium at temperature T . A naive quantum counterpart would be the following master equation [1]:

$$\frac{d\rho}{dt} = -D_{pp}[\mathbf{q}, [\mathbf{q}, \rho]] - i\frac{\eta}{2M}[\mathbf{q}, \{\mathbf{p}, \rho\}] \quad (2)$$

where $\mathbf{q}, \mathbf{p}$ denote the position and, resp., the momentum *operators* in interaction picture. This equation has been used recently to describe the paradigm of environmental decoherence [2]. Eq. (2), however, does not belong to the Lindblad class [3] and, consequently, may violate the positivity of the *density operator* ρ [4]. A detailed dynamic analysis of the quantum Brownian motion in gas environment is thus needed to derive a suitable quantum evolution equation of Lindblad class.

Let us first consider a simple (frictionless) classical kinetic model. The momentum distribution of a Brownian particle, interacting with the particles of the environment, satisfies the following kinetic equation:

$$\frac{d\rho(\mathbf{p})}{dt} = n_0 \int dE d\Omega_i d\Omega_f k^2 \frac{d\sigma(\theta, E)}{d\Omega_f} \rho^\mathcal{E}(\mathbf{k}_i) (\rho(\mathbf{p} - \mathbf{k}_{fi}) - \rho(\mathbf{p})) \quad (3)$$

where n_0 is the density of environmental particles, $\rho^\mathcal{E}$ is their momentum distribution; $d\sigma$ denotes the differential cross section of their scattering on the Brownian particle, while $\mathbf{k}_{fi} = \mathbf{k}_f - \mathbf{k}_i$ is the difference between the final and initial momenta of the scattered particle. Further symbols are used in a standard way and need no special explanation. The Eq. (3) is valid provided: the typical scattering time is much smaller than the average period between subsequent collisions, the mass M of the Brownian particle is much bigger than the mass m of the environment particles, their interaction is spin independent, and the influence of the Brownian particle on the environment's momentum distribution $\rho^\mathcal{E}$ can be ignored.

Let us illustrate the additional issues arising in the quantum case. Looking for a master equation for the density operator ρ , observe the classical Eq. (3) would govern the evolution of its diagonal elements according to the substitution $\rho(\mathbf{p}) = \langle \mathbf{p} | \rho | \mathbf{p} \rangle$. For the off-diagonal part of ρ , however, we have no evolution rule. Such quantum master equations, constructed after Pauli's pioneering one [5], are nothing to do with quantum states which are not translation invariant. A complete quantum kinetic equation must specify rules for ρ 's off-diagonal part as well. Such an extension of the "Pauli" evolution Eq. (3) would be

$$\frac{d\rho}{dt} = n_0 \int dE d\Omega_i d\Omega_f k^2 \frac{d\sigma(\theta, E)}{d\Omega_f} \rho^\mathcal{E}(\mathbf{k}_i) (e^{-i\mathbf{k}_{fi}\cdot\mathbf{q}} \rho e^{i\mathbf{k}_{fi}\cdot\mathbf{q}} - \rho). \quad (4)$$

This equation was, in fact, derived from the unitary dynamics of the Brownian particle interacting with the gas environment [6,7]).

The quantum master Eq. (4) is not completely realistic since it ignores dissipation (friction). In this Letter we present a systematic derivation of the dissipative quantum master equation. We start from the unitary dynamics of the density operator $\rho^{\mathcal{P}+\mathcal{E}}$ corresponding to the Brownian particle ($\mathcal{P}$) plus the environment ($\mathcal{E}$) coupled to each other. In interaction picture, a single collision between the Brownian and an environmental particle transforms the initial state $\rho_i^{\mathcal{P}+\mathcal{E}}$ into the final one $\rho_f^{\mathcal{P}+\mathcal{E}}$ via the unitary scattering operator S :

$$\rho_f^{\mathcal{P}+\mathcal{E}} = S \rho_i^{\mathcal{P}+\mathcal{E}} S^\dagger. \quad (5)$$

Introduce transition operator T by $S = 1 + iT$. From unitarity of S we get $T^\dagger T = i(T^\dagger - T)$; then Eq. (5) yields [8]:

$$\Delta\rho^{\mathcal{P}+\mathcal{E}} \equiv \rho_f^{\mathcal{P}+\mathcal{E}} - \rho_i^{\mathcal{P}+\mathcal{E}} = \frac{i}{2} [T + T^\dagger, \rho_i^{\mathcal{P}+\mathcal{E}}] + T \rho_i^{\mathcal{P}+\mathcal{E}} T^\dagger - \frac{1}{2} \{T^\dagger T, \rho_i^{\mathcal{P}+\mathcal{E}}\}. \quad (6)$$

The change $\Delta\rho$ of the Brownian particle's density operator is given by the trace of Eq. (6) over the environmental degrees of freedom. We shall approximate $d\rho/dt$ by

$$\frac{\Delta\rho}{\Delta t} = \frac{1}{\Delta t} \text{tr}_\mathcal{E} \Delta\rho^{\mathcal{P}+\mathcal{E}} \quad (7)$$

where Δt is the period spanned by the initial and final states according to Eq. (5), considered longer than the typical collision time.

Consider the standard form of the transition operator for the (spin-independent) scattering of environmental particle on the Brownian one, with initial laboratory momenta $\mathbf{k}_i$ and $\mathbf{p}_i$, resp.:

$$T = \frac{1}{2\pi m^*} \int d\mathbf{p}_i d\mathbf{k}_i d\mathbf{k}_f f(\mathbf{k}_f^*, \mathbf{k}_i^*) \delta(E_{\mathbf{k}_f^*} - E_{\mathbf{k}_i^*}) |\mathbf{p}_i - \mathbf{k}_{fi}, \mathbf{k}_f \rangle \langle \mathbf{p}_i, \mathbf{k}_i|. \quad (8)$$

This equation is valid in laboratory system though c.m.s. quantities (marked by stars) appear in it: $M^* = M + m$, $m^* = mM/M^*$, $\mathbf{k}_i^* = (M/M^*)\mathbf{k}_i - (m/M^*)\mathbf{p}_i$, and $\mathbf{k}_f^* = \mathbf{k}_f - (m/M^*)(\mathbf{k}_i + \mathbf{p}_i)$. Also f is the c.m.s. scattering amplitude.

Assume now the form $\rho_i \otimes \rho^\mathcal{E}$ for $\rho_i^{\mathcal{P}+\mathcal{E}}$. Let the environment's state $\rho^\mathcal{E}$ be stationary, representing identical particles of spatial density n_0 and of identical momentum distribution $\rho^\mathcal{E}(\mathbf{k})$. (Assume Boltzmann statistics, for simplicity's.) Let us substitute Eq. (8) into Eq. (6) and single out the second term on the R.H.S.:

$$\begin{aligned} \text{tr}_\mathcal{E} (T \rho_i \otimes \rho^\mathcal{E} T^\dagger) &= \frac{2\pi n_0}{m^{*2}} \int \int d\mathbf{p}_i d\mathbf{p}'_i d\mathbf{k}_i d\mathbf{k}_f \rho^\mathcal{E}(\mathbf{k}_i) \\ & f(\mathbf{k}_f^*, \mathbf{k}_i^*) \delta(E_{\mathbf{k}_f^*} - E_{\mathbf{k}_i^*}) |\mathbf{p}_i - \mathbf{k}_{fi} \rangle \langle \mathbf{p}_i| \rho |\mathbf{p}'_i \rangle \langle \mathbf{p}'_i - \mathbf{k}_{fi}| f^*(\mathbf{k}_f^{*'}, \mathbf{k}_i^{*'}) \delta(E_{\mathbf{k}_f^{*'}} - E_{\mathbf{k}_i^{*'}}). \end{aligned} \quad (9)$$

Change the integration variables $d\mathbf{k}_i, d\mathbf{k}_f$ for $d\mathbf{k}_i^*, d\mathbf{k}_f^*$ respectively. If $\langle \mathbf{p}_i | \rho | \mathbf{p}'_i \rangle$ is diagonal or *almost* diagonal then, to a good approximation:

$$\begin{aligned} \text{tr}_\mathcal{E} (T \rho \otimes \rho^\mathcal{E} T^\dagger) &= \frac{2\pi n_0}{m^{*2}} \left(\frac{M^*}{M} \right)^3 \int \int d\mathbf{p}_i d\mathbf{p}'_i d\mathbf{k}_i^* d\mathbf{k}_f^* |f(\mathbf{k}_f^*, \mathbf{k}_i^*)|^2 \left[\delta(E_{\mathbf{k}_f^*} - E_{\mathbf{k}_i^*}) \right]^2 \\ & \rho^\mathcal{E}(\mathbf{k}_i) |\mathbf{p}_i - \mathbf{k}_{fi} \rangle \langle \mathbf{p}_i| \rho |\mathbf{p}'_i \rangle \langle \mathbf{p}'_i - \mathbf{k}_{fi}|. \end{aligned} \quad (10)$$

Rewrite it in operator form:

$$\begin{aligned} \text{tr}_\mathcal{E} (T \rho \otimes \rho^\mathcal{E} T^\dagger) &= \frac{2\pi n_0}{m^{*2}} \left(\frac{M^*}{M} \right)^3 \int \int d\mathbf{k}_i^* d\mathbf{k}_f^* |f(\mathbf{k}_f^*, \mathbf{k}_i^*)|^2 \left[\delta(E_{\mathbf{k}_f^*} - E_{\mathbf{k}_i^*}) \right]^2 \\ & \sqrt{\rho^\mathcal{E}(\mathbf{k}_i)} e^{-i\mathbf{k}_{fi}^* \mathbf{q}} \rho e^{i\mathbf{k}_{fi}^* \mathbf{q}} \sqrt{\rho^\mathcal{E}(\mathbf{k}_i)} \end{aligned} \quad (11)$$

where $\mathbf{k}_i = \mathbf{k}_i^* + (m/M)(\mathbf{p} + \mathbf{k}_f^*)$. Apply the usual approximation $\delta(E)|_{E=0} \sim \Delta t/2\pi$. Eq.(11) thus takes the following form:

$$\frac{1}{\Delta t} \text{tr}_\mathcal{E} (T \rho \otimes \rho^\mathcal{E} T^\dagger) = \frac{n_0}{m^{*2}} \left(\frac{M^*}{M} \right)^3 \int \int d\mathbf{k}_i^* d\mathbf{k}_f^* \delta(E_{\mathbf{k}_f^*} - E_{\mathbf{k}_i^*}) |f(\mathbf{k}_f^*, \mathbf{k}_i^*)|^2 V_{\mathbf{k}_f^* \mathbf{k}_i^*} \rho V_{\mathbf{k}_f^* \mathbf{k}_i^*}^\dagger \quad (12)$$

with

$$V_{\mathbf{k}_f^* \mathbf{k}_i^*} = \sqrt{\rho^\mathcal{E} \left(\mathbf{k}_i^* + \frac{m}{M}(\mathbf{p} + \mathbf{k}_f^*) \right)} e^{-i\mathbf{k}_{fi}^* \mathbf{q}}. \quad (13)$$

Now we have completely prepared to calculate the rate $d\rho/dt$ which we approximate by $\Delta\rho/\Delta t$ (7). Consider the R.H.S. of Eq. (6). The contribution of its first term will be neglected (that can be justified in rarified gases), the contribution of the second one is given by Eq. (12), and the third term's yield would also be given by a properly altered form of Eq. (12). Invoking Eq. (7), too, all this reads:

$$\frac{d\rho}{dt} = \frac{n_0}{m^{*2}} \left(\frac{M^*}{M} \right)^3 \int \int d\mathbf{k}_i^* d\mathbf{k}_f^* \delta(E_{\mathbf{k}_f^*} - E_{\mathbf{k}_i^*}) |f(\mathbf{k}_f^*, \mathbf{k}_i^*)|^2 \left(V_{\mathbf{k}_f^* \mathbf{k}_i^*} \rho V_{\mathbf{k}_f^* \mathbf{k}_i^*}^\dagger - \frac{1}{2} \{ V_{\mathbf{k}_f^* \mathbf{k}_i^*}^\dagger V_{\mathbf{k}_f^* \mathbf{k}_i^*}, \rho \} \right). \quad (14)$$

An equivalent equation can be given in terms of the c.m.s. differential cross section $d\sigma/d\Omega = |f|^2$:

$$\frac{d\rho}{dt} = n_0 \left(\frac{M^*}{M} \right)^3 \int dE^* d\Omega_i^* d\Omega_f^* k^{*2} \frac{d\sigma(\theta^*, E^*)}{d\Omega_f^*} \left(V_{\mathbf{k}_f^* \mathbf{k}_i^*} \rho V_{\mathbf{k}_f^* \mathbf{k}_i^*}^\dagger - \frac{1}{2} \{ V_{\mathbf{k}_f^* \mathbf{k}_i^*}^\dagger V_{\mathbf{k}_f^* \mathbf{k}_i^*}, \rho \} \right). \quad (15)$$

This is the central result of our Letter: a quantum master equation of Lindblad form [3] with Lindblad generators $V_{\mathbf{k}_f^* \mathbf{k}_i^*}$, describing the fluctuational-frictional evolution of the Brownian particle's density operator ρ , valid at time scales longer than the typical collision time.

The master Eq. (15) preserves translation invariance: diagonal (in momentum) density operators remain diagonal. Eq.(15) implies the following closed equation for translation invariant states:

$$\frac{d\rho(\mathbf{p})}{dt} = n_0 \left(\frac{M^*}{M} \right)^3 \int dE^* d\Omega_i^* d\Omega_f^* k^{*2} \frac{d\sigma(\theta^*, E^*)}{d\Omega_f^*} \left(\rho^\mathcal{E} \left(\frac{M^*}{M} \mathbf{k}_i^* + \frac{m}{M} (\mathbf{p} - \mathbf{k}_{fi}^*) \right) \rho(\mathbf{p} - \mathbf{k}_{fi}^*) - \rho^\mathcal{E} \left(\frac{M^*}{M} \mathbf{k}_i^* + \frac{m}{M} \mathbf{p} \right) \rho(\mathbf{p}) \right). \quad (16)$$

where, as before, the normalized distribution $\rho(\mathbf{p})$ is defined by $\rho(\mathbf{p}) = \text{const.} \langle \mathbf{p} | \rho | \mathbf{p} \rangle$. This equation can, at the same time, be considered the extension of the classical kinetic Eq. (3) including friction this time. To find the stationary solution of the quantum master equation (15) is easy. Since it must be translationally invariant its diagonal will be subjected to the classical equation (16). Invoking the symmetry of $d\sigma$ for $\mathbf{k}_i^*, \mathbf{k}_f^*$ interchanged, the condition of $d\rho/dt = 0$ will be the following:

$$\rho^\mathcal{E}(\mathbf{k}_i) \rho(\mathbf{p}_i) = \rho^\mathcal{E}(\mathbf{k}_f) \rho(\mathbf{p}_f). \quad (17)$$

i.e. one obtains the *detailed balance* condition. It has very important consequences. Assume the environment is in thermal equilibrium at temperature T , with Boltzmann distribution $\rho^\mathcal{E}(k) = \text{const.} \exp(-\beta E)$ where $\beta \equiv 1/k_B T$. Then Eq. (17) implies that the stationary state ρ of the Brownian particle will be the thermal equilibrium state $\text{const.} \exp(-\beta \mathbf{p}^2/2M)$ in both classical and quantum cases.

Although various special limits of the general master Eq. (15) would be relevant we consider the most traditional one, i.e., the motion of heavy Brownian particle in low density Boltzmann gas, with Ohmic (proportional to velocity) dissipation. In this limiting case $m/M \rightarrow 0$ but $\frac{mp}{Mk}$ is finite. Then Eq.(15) yields:

$$\frac{d\rho}{dt} = n_0 \int dE d\Omega_i d\Omega_f k^2 \frac{d\sigma(\theta, E)}{d\Omega_f} \rho^{\mathcal{E}}(k_i) \left(V_{\mathbf{k}_f \mathbf{k}_i} \rho V_{\mathbf{k}_f \mathbf{k}_i}^\dagger - \frac{1}{2} \{ V_{\mathbf{k}_f \mathbf{k}_i}^\dagger V_{\mathbf{k}_f \mathbf{k}_i}, \rho \} \right). \quad (18)$$

The Lindblad generators (13) will be replaced by:

$$V_{\mathbf{k}_f \mathbf{k}_i} = \left(1 - \frac{\beta}{2M} \mathbf{k}_i \mathbf{p} \right) \exp(-i \mathbf{k}_{fi} \mathbf{q}). \quad (19)$$

The Eq. (18) with the Lindblad generators (19) belong to the class of Ohmic dissipative master equations found phenomenologically by Gallis [9].

Among the early results concerning fenomenological quantum master equations with friction, Dekker's proposal [10] is worth to mention. It is recovered by our Eqs. (18,19) in the limit when coherent extension of the Brownian particle and/or transferred momentum $|\mathbf{k}_{fi}|$ in single collisions are small enough. Then $\exp(-i \mathbf{k}_{fi} \mathbf{q})$ on the R.H.S. of Eq. (19) is approximated by $1 - i \mathbf{k}_{fi} \mathbf{q}$ and, neglecting terms of order of $q^2 p^2$, Eq. (18) can be written as follows:

$$\begin{aligned} \frac{d\rho}{dt} = n_0 \int dE d\Omega_i d\Omega_f k^2 \frac{d\sigma(\theta, E)}{d\Omega_f} \rho^{\mathcal{E}}(k_i) & \left(i \frac{\beta}{4M} [\{\mathbf{k}_i \mathbf{p}, \mathbf{k}_{fi} \mathbf{q}\}, \rho] \right. \\ & \left. + (\mathbf{k}_{fi} \mathbf{q} - i \frac{\beta}{2M} \mathbf{k}_i \mathbf{p}) \rho (\mathbf{k}_{fi} \mathbf{q} + i \frac{\beta}{2M} \mathbf{k}_i \mathbf{p}) - \frac{1}{2} \{ (\mathbf{k}_{fi} \mathbf{q} + i \frac{\beta}{2M} \mathbf{k}_i \mathbf{p}) (\mathbf{k}_{fi} \mathbf{q} - i \frac{\beta}{2M} \mathbf{k}_i \mathbf{p}), \rho \} \right). \end{aligned} \quad (20)$$

This is identical to Dekker's fenomenological equation (in interaction picture):

$$\frac{d\rho}{dt} = -D_{pp}[\mathbf{q}, [\mathbf{q}, \rho]] - D_{qq}[\mathbf{p}, [\mathbf{p}, \rho]] - i \frac{\eta}{2M} [\mathbf{q}, \{\mathbf{p}, \rho\}]. \quad (21)$$

We can express Dekker's parameters as follows:

$$D_{pp} = \frac{4}{3} n_0 \int dE d\Omega_i d\Omega_f k^4 \frac{d\sigma(\theta, E)}{d\Omega_f} \rho^{\mathcal{E}}(k_i) \sin^2 \frac{\theta}{2}, \quad (22)$$

$$D_{qq} = \frac{1}{3} n_0 \left(\frac{\beta}{2M} \right)^2 \int dE d\Omega_i d\Omega_f k^4 \frac{d\sigma(\theta, E)}{d\Omega_f} \rho^{\mathcal{E}}(k_i), \quad (23)$$

$$\eta = \beta D_{pp}. \quad (24)$$

One can now compare the naive quantum master equation (2) to our result (21-24): we have got an additional term of *position diffusion* definitely absent in the classical equation (1). Classical kinematics, even if stochastic, is based on continuous trajectories in configurational space and does not allow position diffusion at all since this latter would assume finite position jumps. At the same time, momentum jumps are reasonable idealizations in classical mechanics. Let us resume a particular lesson of the longstanding

controversies related to the "true" quantum counterpart of the classical Brownian motion: quantum friction is accompanied by momentum *and position* diffusions, which latter is neither present nor even possible in classical kinematics nonetheless without which no mathematically consistent (i.e. Lindblad) quantum master equation would be written down [11].

This work was supported by OTKA Grant No.1822/1991.

^(a)Electronic address: diosi@rmki.kfki.hu

- [1] A. O. Caldeira and A. J. Leggett, *Physica* **121A**, 587 (1983).
- [2] W. H. Zurek, *Phys. Today* **44**, No. 10, 36 (1991).
- [3] G. Lindblad, *Commun. Math. Phys.* **48**, 119 (1976).
- [4] V. Ambegaokar, *Ber. Bunsenes. Phys. Chem.* **95**, 400 (1991).
- [5] W. Pauli, *Collected Scientific Papers* (Interscience, New York, 1964).
- [6] E. Joos and H. D. Zeh, *Z. Phys.* **B59**, 223 (1985).
- [7] M. R. Gallis, *Phys. Rev.* **42A**, 38 (1990).
- [8] S. Stenholm, *Phys.Scri.* **47**, 724 (1993)
- [9] M. R. Gallis, *Phys. Rev.* **48A**, 1028 (1993).
- [10] H. Dekker, *Phys. Rep.* **80**, 1 (1981).
- [11] L. Diósi, *Europhys. Lett.* **22**, 1 (1993); *Physica*, 199A, 517 (1993).